

Medications and Birth-Control Pills

Several studies have shown that taking these medications may reduce the effectiveness of your birth-control pills:

antibiotics — sulfa drugs
pain killers — tranquilizers
sedatives or antihistamines

Please check with your doctor before taking these or any other medication.

Medications and Birth-Control Pills

Several studies have shown that taking these medications may reduce the effectiveness of your birth-control pills:

antibiotics — sulfa drugs
pain killers — tranquilizers
sedatives or antihistamines

Please check with your doctor before taking these or any other medication.

Medications and Birth-Control Pills

Several studies have shown that taking these medications may reduce the effectiveness of your birth-control pills:

antibiotics — sulfa drugs
pain killers — tranquilizers
sedatives or antihistamines

Please check with your doctor before taking these or any other medication.

Medications and Birth-Control Pills

Several studies have shown that taking these medications may reduce the effectiveness of your birth-control pills:

antibiotics — sulfa drugs
pain killers — tranquilizers
sedatives or antihistamines

Please check with your doctor before taking these or any other medication.

Medications and Birth-Control Pills

Several studies have shown that taking these medications may reduce the effectiveness of your birth-control pills:

antibiotics — sulfa drugs

pain killers — tranquilizers

sedatives or antihistamines

Please check with your doctor before taking these or any other medication.